

Equivariant heat invariants of the Laplacian and nonminimal operators on differential forms

Yong Wang

Abstract

In this paper, we compute the first two equivariant heat kernel coefficients of the Bochner Laplacian on differential forms. The first two equivariant heat kernel coefficients of the Bochner Laplacian with torsion are also given. We also study the equivariant heat kernel coefficients of nonminimal operators on differential forms and get the equivariant Gilkey-Branson-Fulling formula.

MSC: 58J05, 54A14

Keywords: Equivariant heat kernel asymptotics; Bochner Laplacian; nonminimal operators; Gilkey-Branson-Fulling formula

1 Introduction

In [Do1], Donnelly computed heat kernel coefficients of the Bochner Laplacian with torsion on differential forms. In [Do2], the equivariant heat kernel asymptotics of the Laplacian on functions was established and the first two equivariant heat kernel coefficients were evaluated. Donnelly's results were used to study the asymptotic expansion of the heat kernel for orbifolds in [DGGW]. On the other hand, in [GBF], Gilkey, Branson and Fulling studied the heat equation asymptotics of nonminimal operators on differential forms and computed the first two heat kernel coefficients. In [AV], the heat kernel expansion for a general nonminimal operator on the spaces $C^\infty(\wedge^k)$ and $C^\infty(\wedge^{p,q})$ was studied. In [Gi], Gilkey studied the heat kernel asymptotics of nonminimal operators for manifolds with boundary. In [PC], the Gilkey-Branson-Fulling formula was generalized to the case of the h -Laplacian. The purpose of this paper is to compute the first two equivariant heat kernel coefficients of the Bochner Laplacian and the Bochner Laplacian with torsion on differential forms and give the equivariant version of the Gilkey-Branson-Fulling formula. We firstly compute the first two equivariant heat kernel coefficients of the Bochner Laplacian on differential forms which are given in Section 2. In section 3, we give the first two equivariant heat kernel coefficients of the Bochner Laplacian with torsion on differential forms. In Section 4, we prove the equivariant Gilkey-Branson-Fulling formula.

2 The computation of the equivariant heat kernel coefficients

Let M be a compact oriented Riemannian manifold of dimension d and $T : M \rightarrow M$ be an isometry preserving orientation with fixed point set Ω . Ω is the disjoint union of closed connected submanifolds N of dimension n . Let TM and T^*M be the tangent and cotangent bundles and let $\wedge^p$ be the bundle of exterior p -forms over M and Δ be the Bochner Laplacian associated to the Levi-Civita connection on $\wedge^p$. T induces a map T^* on $\wedge^p$ which commutes with Δ . If λ is an eigenvalue of Δ , then T^* maps the λ eigenspace into itself. We shall be interested in the sum $\sum_{\lambda} \text{Tr}(T_{\lambda}^*)e^{-t\lambda}$, $t > 0$ where Tr denotes the trace. Using the same discussions of Theorem 4.1 in [Do] (also see [Gil]), we get

Theorem 1 *Let $T : M \rightarrow M$ be an isometry with fixed point set Ω . Then there is an asymptotic expansion*

$$\sum_{\lambda} \text{Tr}(T_{\lambda}^*)e^{-t\lambda} \sim \sum_{N \in \Omega} (4\pi t)^{-\frac{n}{2}} \sum_{k=0}^{\infty} t^k \int_N b_k(T, a) d\text{vol}_N(a), \quad t \rightarrow 0, \quad (2.1)$$

where the functions $b_k(T, a)$ depend only upon the germ of T and the Riemannian metric of M near points $a \in N$.

Let $u_i(x, y)$ be the i -th term of the asymptotic expansion of the heat kernel of $e^{-t\Delta}$ and A denote the endomorphism induced by T on the fibre of the normal bundle over a and $B = (I - A)^{-1}$. Let U_N be a sufficient small tubular neighborhood of N and $\pi : U_N \rightarrow N$ be the projection. Let x be coordinates for a normal coordinate chart on $\pi^{-1}(a)$ for $a \in N$ and $\bar{x} = x - T(x)$. Let $d\text{vol}_M(x) = \psi(x)dx(\pi^*d\text{vol}_N(a))$. By the Morse lemma, one can find a smooth coordinate change so that (see Appendix in [Do])

$$d^2(\bar{x} + T(x), T(x)) = \sum_{i=1}^s y_i^2 = |y|^2.$$

Let $|J(\bar{x}, y)|$ denote the absolute value of the Jacobian determinant of this change of variables. Let $\square_y = \sum_{i=1}^s \frac{\partial^2}{\partial y_i^2}$ and

$$h_i(x(y)) = |\det B| \text{tr}[T^* u_i(T(x), x)] |J(\bar{x}, y)| \psi(x), \quad (2.2)$$

then by the same discussions as in [Do], we have

$$b_k(T, a) = \sum_{j=0}^k \frac{1}{j!} \square_y^j (h_{k-j})(0). \quad (2.3)$$

Let $\tau(y, x) : \wedge_x^p \rightarrow \wedge_y^p$ be the parallel transport along the geodesic curve from x to y . Fix x and suppose y is in some normal coordinate neighborhood w_j of x and

$g_{ij} = g(\partial/\partial w_i, \partial/\partial w_j)$. By Chapter 2 in [BGV], we have $u_0(y, x) = (\det g_{ij})^{-\frac{1}{4}} \tau(y, x)$. By (2.3),

$$b_0(T, a) = |\det B| \operatorname{tr}[T^* u_0(T(a), a)] = |\det B| \operatorname{tr}[T_a^*|_{\wedge^p}].$$

We write $\tilde{A} = \begin{pmatrix} I & 0 \\ 0 & A^t \end{pmatrix}$ and define

$$\wedge^p \tilde{A}(\theta_1 \wedge \cdots \wedge \theta_p) = \tilde{A}\theta_1 \wedge \cdots \wedge \tilde{A}\theta_p, \quad (2.4)$$

then

$$\operatorname{tr}[\wedge^p \tilde{A}] = \sum_{0 \leq p_1 \leq p} \sum_{n+1 \leq i_1 < \cdots < i_{p_1} \leq d} \sum_{l_1, \dots, l_{p_1}} \varepsilon_{i_1, \dots, i_{p_1}}^{l_1, \dots, l_{p_1}} A_{i_1 l_1} \cdots A_{i_{p_1} l_{p_1}} \quad (2.5),$$

where $\varepsilon_{i_1, \dots, i_{p_1}}^{l_1, \dots, l_{p_1}}$ is the generalized Kronecker symbol and

$$b_0(T, a) = |\det B| \operatorname{tr}[\wedge^p \tilde{A}]. \quad (2.6)$$

By (2.3),

$$b_1(T, a) = |\det B| (h_1(0) + \square_y (\operatorname{tr}[T^* u_0(T(x), x)] |J(\bar{x}, y)|\psi(x))(0)). \quad (2.7)$$

Denote $\tau_0 = \sum_{a,b=1}^d R_{abab}$ and $\rho_{ab} = \sum_{c=1}^d R_{acbc}$ the scalar curvature and Ricci tensor of M . By Lemma 4.8.7 in [Gil],

$$u_1(a, a) = \frac{\tau_0}{6}, \quad (2.8)$$

so

$$h_1(0) = \operatorname{tr}[T_a^* u_1(T(a), a)] = \frac{\tau_0}{6} \operatorname{tr}[\wedge^p \tilde{A}], \quad (2.9)$$

By the Taylor expansions in page 169 in [Do], we know that

$$\frac{\partial}{\partial y_i} \Big|_{y=0} \left((\det g_{ij})^{-\frac{1}{4}} |J(\bar{x}, y)|\psi(x) \right) = 0, \quad (2.10)$$

so

$$\begin{aligned} \square_y (\operatorname{tr}[T^* u_0(T(x), x)] |J(\bar{x}, y)|\psi(x))(0) &= \square_y (\operatorname{tr}[T^* \tau(T(x), x)])|_{y=0} \\ &+ \operatorname{tr}[\wedge^p \tilde{A}] \square_y \left((\det g_{ij})^{-\frac{1}{4}} |J(\bar{x}, y)|\psi(x) \right) (0). \end{aligned} \quad (2.11)$$

In the following, we adopt the convention of summing Greek indices $1 \leq \alpha, \beta, \gamma \leq n$ from 1 to n and Latin indices $n+1 \leq i, j, k \leq d$ from $n+1$ to d . By the computations in [Do, p.170], we know that

$$\square_y \left((\det g_{ij})^{-\frac{1}{4}} |J(\bar{x}, y)|\psi(x) \right) (0) = \frac{1}{6} \rho_{kk} + \frac{1}{3} R_{iksh} B_{ki} B_{hs} + \frac{1}{3} R_{ikth} B_{kt} B_{hi} - R_{k\alpha h\alpha} B_{ks} B_{hs}. \quad (2.12)$$

Now we compute $\operatorname{tr}[T^* \tau(T(x), x)]$. Let $E = (E_1, \dots, E_d)$ be an oriented orthonormal frame field in a neighborhood of a such that for $x \in N$, $E_1(x), \dots, E_n(x)$ are tangent to N while the vector fields $E_{n+1}(x), \dots, E_d(x)$ are normal to N and E is parallel

along geodesics normal to N and dT is expressed as a matrix-valued function $A(x)$ as $dTE(x) = E(Tx)A(x)$. Then $A(a) = \begin{pmatrix} I & 0 \\ 0 & A \end{pmatrix}$. Let $x = \exp_a \left(\sum_{i=n+1}^d x_i E_i(a) \right)$. By Lemma 3.1 in [LYZ], $A(x) = A(a)$. We consider $\wedge^p(T^*M)$ as the associated bundle $SO(T^*M) \times_\mu \wedge^p(\mathbf{R}^n)$ where the action μ is defined by (2.4). Let $\sigma = (E_1^*, \dots, E_d^*)$ be the local section of $SO(T^*M)$ on V . Then a local section of $\wedge^p(T^*M)$ on V can be expressed as $[(\sigma, f)]$ where $f : V \rightarrow \wedge^p(\mathbf{R}^n)$ be a smooth function. Let

$$T^*[(\sigma(Tx), c)] = [(\sigma(x), \bar{T}^*(x)c)], \quad (2.13)$$

where $\bar{T}^*(x) : V \rightarrow \text{End}(\mathbf{R}^n)$. Let

$$\tau(Tx, x)[(\sigma(x), c)] = [(\sigma(Tx), \bar{\tau}^*(x)c)]. \quad (2.14)$$

Then $\bar{T}^*(x) = \wedge^p \tilde{A}$ is a constant matrix-valued function. As in [LYZ, p.575], define the oriented frame field E^{Tx} over the patch V by requiring that $E^{Tx}(Tx) = E(Tx)$ and that E^{Tx} be parallel along geodesics through Tx and a map $\Phi : V \rightarrow so(d)$ by $E^{Tx}(x) = E(x)e^{\Phi(x)}$. Then $E^{*,x}(Tx) = E^*(Tx)e^{-\Phi(x)}$, that is

$$\tau(Tx, x)\sigma(x) = \sigma(Tx)e^{-\Phi(x)}, \quad (2.15)$$

so $\bar{\tau}^*(x) = \wedge^p e^{-\Phi(x)}$. By Lemma 3.3 in [LYZ], $\Phi = \begin{pmatrix} 0 & 0 \\ 0 & \Psi \end{pmatrix}$ and

$$\Psi_{ij}(x) = -\frac{1}{2}A_{rl}x_l x_s R_{rsij}(a) + O(|x|^3),$$

By $x_i = B_{ij}\bar{x}_j$ and $\bar{x}_j = y_j + O(|y|^3)$, so

$$\begin{aligned} \Psi_{ij}(x) &= -\frac{1}{2}A_{rl}B_{lk}B_{sq}R_{rsij}(a)y_k y_q + O(|y|^3). \\ e^{-\Phi(x)} &= 1 - \Phi(x) + O(|x|^3). \\ \text{tr}[T^*\tau(T(x), x)] &= \text{tr}[\wedge^p(W)] = \text{tr}[\wedge^p(\tilde{A}(1 - \Phi(x)))] + O(|x|^3), \quad (2.16) \\ C &:= \square_y(\text{tr}[\wedge^p(W)])|_{y=0} = \sum_{\delta=1}^{d-n} \sum_{1 \leq i_1 < \dots < i_p \leq d} \sum_{l_1, \dots, l_p} \varepsilon_{i_1, \dots, i_p}^{l_1, \dots, l_p} \sum_{s=1}^p W_{l_1 i_1} \cdots W_{l_{s-1} i_{s-1}} \\ &\quad \cdot \left(\frac{\partial^2}{\partial y_\delta^2} W_{l_s i_s} \right) W_{l_{s+1} i_{s+1}} \cdots W_{l_p i_p} |_{y=0} \\ &= \sum_{\delta=1}^{d-n} \sum_{1 \leq i_1 < \dots < i_p \leq d} \sum_{l_1, \dots, l_p} \varepsilon_{i_1, \dots, i_p}^{l_1, \dots, l_p} \sum_{s=1}^p A_{i_1 l_1} \cdots A_{i_{s-1} l_{s-1}} A_{i_{s+1} l_{s+1}} \cdots A_{i_p l_p} A_{rl} B_{l\delta} B_{v\delta} A_{ml_s} R_{rvmi_s} \\ &= \sum_{\delta=1}^{d-n} \sum_{1 \leq p_1 \leq p} \sum_{n+1 \leq i_1 < \dots < i_{p_1} \leq d} \sum_{l_1, \dots, l_{p_1}} \varepsilon_{i_1, \dots, i_{p_1}}^{l_1, \dots, l_{p_1}} \sum_{s=1}^{p_1} A_{i_1 l_1} \cdots A_{i_{s-1} l_{s-1}} \end{aligned}$$

$$\cdot A_{i_{s+1}l_{s+1}} \cdots A_{i_{p_1}l_{p_1}} A_{rl} B_{l\delta} B_{v\delta} A_{ml_s} R_{vmi_s}. \quad (2.17)$$

So by (2.5),(2.7),(2.11),(2.12) and (2.17), we get

$$\begin{aligned} b_1(T, a) = |\det B| \left\{ C + \operatorname{tr}[\wedge^p \tilde{A}] \left(\frac{\tau_0}{6} + \frac{1}{6} \rho_{kk} \right. \right. \\ \left. \left. + \frac{1}{3} R_{iksh} B_{ki} B_{hs} + \frac{1}{3} R_{ikth} B_{kt} B_{hi} - R_{k\alpha h\alpha} B_{ks} B_{hs} \right) \right\}. \end{aligned} \quad (2.18)$$

Theorem 2 *The coefficient $b_k(T, a)$ of Theorem 1 is of the form $b_k(T, a) = |\det B| b'_k(T, a)$ where $b'_k(T, a)$ is an invariant polynomial in the components of A , B and the curvature tensor R and its covariant derivative at a . In particular, $b_0(T, a)$, $b_1(T, a)$ are determined by (2.6) and (2.18).*

Remark Theorem 2 can be used to evaluate the heat kernel coefficients of the Laplacian on forms on orbifolds as in [DGGW].

3 The computation of the equivariant heat kernel coefficients of the Bochner Laplacian with torsion

The Levi-Civita connection ∇ is the unique torsion zero connection on TM which preserves the metric. More generally, let $\overline{T} : TM \times TM \rightarrow TM$ be a skew-symmetric linear map, i.e. $\overline{T}(X, Y) = -\overline{T}(Y, X)$. Then there is exactly one metric-preserving connection $\overline{\nabla} = \nabla + Q$ on TM with torsion tensor $\overline{T}$. Let $Q(X, Y, Z) = g(Q(X, Y), Z)$; $\overline{T}(X, Y, Z) = g(\overline{T}(X, Y), Z)$ and $Q_{ijk} = Q(E_i, E_j, E_k)$; $\overline{T}_{ijk} = \overline{T}(E_i, E_j, E_k)$, then

$$Q_{ijk} = \frac{1}{2}(\overline{T}_{ijk} + \overline{T}_{kij} + \overline{T}_{kji}). \quad (3.1)$$

Let $\overline{\nabla}^* = \nabla^* + Q^*$ be the dual connection of $\overline{\nabla}$ on T^*M and $\overline{Q}_{ijk} = g(Q^*(E_i)E_j^*, E_k^*)$, then $\overline{Q}_{ikj} = -Q_{ijk}$. $\overline{\nabla}^*$ induces a connection on $\wedge^p T^*M$. We still denote it by $\overline{\nabla}^*$. Let the Bochner Laplacian with torsion be

$$\overline{\Delta} = - \sum_{i=1}^d (\overline{\nabla}_{E_i}^* \overline{\nabla}_{E_i}^* - \overline{\nabla}_{\nabla_{E_i}^L E_i}^*).$$

Let $\overline{\tau}(y, x) : \wedge_x^p \rightarrow \wedge_y^p$ be the parallel transport along the geodesic curve from x to y associated to the connection $\overline{\nabla}^*$. Define the oriented frame field $\overline{E}^{x,*}$ over the patch V by requiring that $\overline{E}^{x,*}(x) = E^*(x)$ and that $\overline{E}^{x,*}$ be parallel along geodesics through x associated to the connection $\overline{\nabla}^*$. Let z be the normal coordinates for the center x . Let

$$\overline{E}^{x,*}(z) = E^{x,*}(z) \overline{\Psi}(x, z); \quad \overline{\Psi}(x, Tx) = \overline{\Psi}(x). \quad (3.2)$$

Nextly, we compute the Taylor expansion of $\overline{\Psi}(x)$. Let $Q^*E^{x,*}(z) = E^{x,*}(z)A^x(z)$ and $L(x) = A^x(0)$, then

$$Q^*E^*(x) = E^*(x)L(x), \quad L(E_k)_{ij} = Q_{kji}. \quad (3.3)$$

Let γ be the geodesic curve from x to z . By $\nabla_{\dot{\gamma}}^{L,*}E^{x,*} = 0$, $\overline{\nabla}_{\dot{\gamma}}^*E^{x,*} = 0$ and (3.2), we have

$$\frac{d}{dt}\overline{\Psi}(x, \gamma(t)) + A^x(\dot{\gamma}(t))\overline{\Psi}(x, \gamma(t)) = 0. \quad (3.4)$$

By $\gamma(t) = tz$; $\dot{\gamma} = \sum_{j=1}^d z_j \frac{\partial}{\partial z_j}$,

$$\sum_{j=1}^d \frac{\partial \overline{\Psi}}{\partial z_j}(x, tz) z_j + \sum_{j=1}^d z_j A^x\left(\frac{\partial}{\partial z_j}\right)(tz) \overline{\Psi}(x, tz) = 0. \quad (3.5)$$

By $\overline{\Psi}(x, 0) = Id$ and setting $t = 0$, we get

$$\sum_{j=1}^d \frac{\partial \overline{\Psi}}{\partial z_j}(x, 0) z_j + \sum_{j=1}^d z_j A^x\left(\frac{\partial}{\partial z_j}\right)(0) = 0. \quad (3.6)$$

Taking the derivative about t and setting $t = 0$, we get

$$\sum_{i,j=1}^d \frac{\partial^2 \overline{\Psi}}{\partial z_i \partial z_j}(x, 0) z_i z_j + \sum_{i,j=1}^d z_i z_j A^x\left(\frac{\partial}{\partial z_j}\right)(0) \frac{\partial \overline{\Psi}}{\partial z_i}(x, 0) + \sum_{i,j=1}^d z_i z_j \frac{\partial A^x(\frac{\partial}{\partial z_j})}{\partial z_i}(0) = 0. \quad (3.7)$$

By (3.6) and (3.7), we have

$$\begin{aligned} \overline{\Psi}(x, z) &= Id - \sum_{j=1}^d z_j A^x\left(\frac{\partial}{\partial z_j}\right)(0) + \frac{1}{2} \sum_{i,j=1}^d A^x\left(\frac{\partial}{\partial z_j}\right)(0) A^x\left(\frac{\partial}{\partial z_i}\right)(0) z_i z_j \\ &\quad - \frac{1}{2} \sum_{i,j=1}^d z_i z_j \frac{\partial A^x(\frac{\partial}{\partial z_j})}{\partial z_i}(0) + O(|z|^3). \end{aligned} \quad (3.8)$$

Let $Tx = \exp_x(\sum_{i=1}^d u_i E_i(x))$ and $x = (a, c)$ be the orthogonal coordinates in [LYZ, p.574], then by the proposition in [Yu, p.84], we have

$$u_i = O(|c|^3), \quad 1 \leq i \leq n, \quad u_i = \overline{c_{i-n}} - c_{i-n} + O(|c|^3), \quad n+1 \leq i \leq d, \quad (3.9)$$

where $c = (\overline{c}_1, \dots, \overline{c}_{d-n})A$. Then

$$\begin{aligned} \overline{\Psi}(x) &= \overline{\Psi}(x, u) = Id - \sum_{j=n+1}^d u_j A^x\left(\frac{\partial}{\partial z_j}\right)(0) + \frac{1}{2} \sum_{i,j=n+1}^d A^x\left(\frac{\partial}{\partial z_j}\right)(0) A^x\left(\frac{\partial}{\partial z_i}\right)(0) u_i u_j \\ &\quad - \frac{1}{2} \sum_{i,j=n+1}^d u_i u_j \frac{\partial A^x(\frac{\partial}{\partial z_j})}{\partial z_i}(0) + O(|u|^3). \end{aligned} \quad (3.10)$$

We know that

$$A^x\left(\frac{\partial}{\partial z_j}\right)(0) = A^x(0)(E_j(x)) = L(x)(E_j(x)), \quad (3.11)$$

$$\frac{\partial A^{(a,0)}\left(\frac{\partial}{\partial z_j}\right)}{\partial z_j}(0) = \frac{\partial A^{(a,0)}\left(\frac{\partial}{\partial c_{j-n}}\right)}{\partial c_{j-n}}(0) = \frac{\partial L(E_j)}{c_{j-n}}(0), \quad n+1 \leq j \leq d, \quad (3.12)$$

$$L(E_j) = L(E_j)(a, 0) + \sum_{i=n+1}^d \frac{\partial L(E_j)}{\partial c_{i-n}}(a, 0)c_{i-n} + O(|c|^2), \quad (3.13)$$

so

$$\begin{aligned} \overline{\Psi}(x) &= Id - \sum_{j=n+1}^d u_j L(E_j)(a, 0) - \sum_{i,j=n+1} \frac{\partial L(E_j)}{\partial c_{i-n}}(a, 0)c_{i-n}u_j \\ &+ \frac{1}{2} \sum_{i,j=n+1}^d L(E_i)(a, 0)L(E_j)(a, 0)u_i u_j - \frac{1}{2} \sum_{i,j=n+1} \frac{\partial L(E_j)}{\partial c_{i-n}}(a, 0)u_i u_j + O(|c|^3). \end{aligned} \quad (3.14)$$

By

$$u_i = -y_{i-n} + O(|y|^3), \quad c_{i-n} = B_{ij}y_{j-n} + O(|y|^3), \quad n+1 \leq i, j \leq d, \quad (3.15)$$

we have

$$\begin{aligned} \overline{\Psi}(x) &= Id + \sum_{j=n+1}^d y_{j-n} L(E_j)(a, 0) + \sum_{i,j=n+1} \frac{\partial L(E_j)}{\partial c_{i-n}}(a, 0)y_{j-n}B_{ik}y_{k-n} \\ &+ \frac{1}{2} \sum_{i,j=n+1}^d L(E_i)(a, 0)L(E_j)(a, 0)y_{i-n}y_{j-n} - \frac{1}{2} \sum_{i,j=n+1} \frac{\partial L(E_j)}{\partial c_{i-n}}(a, 0)y_{i-n}y_{j-n} + O(|y|^3). \end{aligned} \quad (3.16)$$

By (3.2),

$$\overline{E}^{x,*}(Tx) = E^*(Tx)e^{-\Phi(x)}\overline{\Psi}(x). \quad (3.17)$$

Let

$$\overline{\tau}(Tx, x)[(\sigma(x), c)] = [(\sigma(Tx), \tilde{\tau}^*(x)c)]. \quad (3.18)$$

Let $\overline{b}_i(T, a)$ denote the equivariant heat kernel coefficients of the Bochner Laplacian with torsion. Similar to the discussions in Section 2, $\overline{b}_0(T, a) = b_0(T, a)$. Similar to (2.18), we have

$$\begin{aligned} \overline{b}_1(T, a) &= |\det B| \left\{ \overline{C} + \text{tr}[\wedge^p \tilde{A}]\left(\frac{\tau_0}{6} + \frac{1}{6}\rho_{kk}\right) \right. \\ &\quad \left. + \frac{1}{3}R_{iksh}B_{ki}B_{hs} + \frac{1}{3}R_{ikth}B_{kt}B_{hi} - R_{k\alpha h\alpha}B_{ks}B_{hs} \right\}. \end{aligned} \quad (3.19)$$

where

$$\overline{C} = \square_y(\text{tr}[T^*\tilde{\tau}(T(x), x)])|_{y=0} = \text{tr}[\wedge^p(\tilde{A}e^{-\Phi(x)}\overline{\Psi}(x))] := \text{tr}[\wedge^p \overline{W}]. \quad (3.20)$$

So

$$\begin{aligned}\overline{C} &= \sum_{\delta=1}^{d-n} \sum_{1 \leq i_1 < \dots < i_p \leq d} \sum_{l_1, \dots, l_p} \varepsilon_{i_1, \dots, i_p}^{l_1, \dots, l_p} \frac{\partial^2}{\partial y_\delta^2} [\overline{W}_{l_1 i_1} \dots \overline{W}_{l_p i_p}]|_{y=0} \\ &= \sum_{\delta=1}^{d-n} \sum_{1 \leq i_1 < \dots < i_p \leq d} \sum_{l_1, \dots, l_p} \varepsilon_{i_1, \dots, i_p}^{l_1, \dots, l_p} \left[2 \sum_{1 \leq m_1 < m_2 \leq p} \overline{W}_{l_1 i_1} \dots \frac{\partial \overline{W}_{l_{m_1} i_{m_1}}}{\partial y_\delta} \dots \frac{\partial \overline{W}_{l_{m_2} i_{m_2}}}{\partial y_\delta} \dots \overline{W}_{l_p i_p} \right. \\ &\quad \left. \sum_{1 \leq m_3 \leq p} \overline{W}_{l_1 i_1} \dots \frac{\partial^2 \overline{W}_{l_{m_3} i_{m_3}}}{\partial y_\delta^2} \dots \overline{W}_{l_p i_p} \right] |_{y=0}\end{aligned}$$

By (3.3) and (3.16)

$$\begin{aligned}\frac{\partial \overline{\Psi}_{\varepsilon i_{m_1}}}{\partial y_\delta} |_{y=0} &= -Q_{\delta+n} \varepsilon i_{m_1}; \\ \frac{\partial^2 \overline{\Psi}_{a,b}}{\partial y_\delta^2} |_{y=0} &= 2 \sum_{j=n+1}^d Q_{\delta+n} b_{a,j} B_{j\delta+n} + \sum_{a_1=1}^d Q_{\delta+n} a_1 a Q_{\delta+n} b_{a_1} - Q_{\delta+n} b_{a,\delta+n}, \quad (3.21)\end{aligned}$$

then

$$\begin{aligned}\frac{\partial \overline{W}_{l_{m_1} i_{m_1}}}{\partial y_\delta} |_{y=0} &= -A_{\varepsilon l_{m_1}} Q_{\delta+n} \varepsilon i_{m_1}; \quad (3.22) \\ \frac{\partial^2 \overline{W}_{l_{m_3} i_{m_3}}}{\partial y_\delta^2} |_{y=0} &= A_{\varepsilon_1 l_{m_3}} [A_{rl} B_{l\delta} B_{v\delta} A_{m\varepsilon_1} R_{rvmi_{m_3}} + 2 \sum_{j=n+1}^d Q_{\delta+n} i_{m_3} \varepsilon_1, j B_{j\delta+n} \\ &\quad + \sum_{a_1=1}^d Q_{\delta+n} a_1 \varepsilon_1 Q_{\delta+n} i_{m_3} a_1 - Q_{\delta+n} i_{m_3} \varepsilon_1, \delta+n]. \quad (3.23)\end{aligned}$$

So

$$\begin{aligned}\overline{C} &= \sum_{\delta=1}^{d-n} \sum_{1 \leq i_1 < \dots < i_p \leq d} \sum_{l_1, \dots, l_p} \varepsilon_{i_1, \dots, i_p}^{l_1, \dots, l_p} \left[2 \sum_{1 \leq m_1 < m_2 \leq p} A_{l_1 i_1} \dots \hat{A}_{l_{m_1} i_{m_1}} \dots \hat{A}_{l_{m_2} i_{m_2}} \dots A_{l_p i_p} \right. \\ &\quad \cdot A_{\varepsilon_1 l_{m_1}} Q_{\delta+n} i_{m_1} \varepsilon_1 A_{\varepsilon_2 l_{m_2}} Q_{\delta+n} i_{m_2} \varepsilon_2 + \sum_{1 \leq m_3 \leq p} A_{l_1 i_1} \dots \hat{A}_{l_{m_3} i_{m_3}} \dots A_{l_p i_p} \\ &\quad \cdot A_{\varepsilon_1 l_{m_3}} [A_{rl} B_{l\delta} B_{v\delta} A_{m\varepsilon_1} R_{rvmi_{m_3}} + 2 \sum_{j=n+1}^d Q_{\delta+n} i_{m_3} \varepsilon_1, j B_{j\delta+n} \\ &\quad \left. + \sum_{a_1=1}^d Q_{\delta+n} a_1 \varepsilon_1 Q_{\delta+n} i_{m_3} a_1 - Q_{\delta+n} i_{m_3} \varepsilon_1, \delta+n] \right] \quad (3.24)\end{aligned}$$

Theorem 3 The coefficient $\overline{b}_k(T, a)$ is of the form $\overline{b}_k(T, a) = |\det B| \overline{b}_k(T, a)$ where $\overline{b}_k(T, a)$ is an invariant polynomial in the components of A , B and the curvature tensor R and the torsion tensor $\overline{T}$ and its covariant derivative at a . In particular, $\overline{b}_1(T, a)$ are determined by (3.24) and (3.19).

In the following, we define another Bochner's Laplacian with torsion and compute its equivariant heat invariants. Let

$$\widehat{\Delta} = - \sum_{i=1}^d (\nabla_{E_i}^* \nabla_{E_i}^* - \nabla_{\bar{\nabla}_{E_i} E_i}^*) = \Delta + Q_{ij} E_j + F.$$

Let $\widehat{\nabla} = \nabla - \frac{1}{2}(Q(E_i)E_i)^*$ be a connection on $\wedge T^p M$ associated to a connection $\widehat{\nabla} = \nabla - \frac{1}{2p}(Q(E_i)E_i)^*$ on T^*M . Then $\widehat{\Delta}$ is a generalized Laplacian associated to the connection $\widehat{\nabla}$. In this case, $L(x) = -\frac{1}{2p}(Q(E_i)E_i)^*$ and $L(E_j) = -\frac{1}{2p}Q_{ij}$. By Proposition 3.3, 3.5 and 4.3 [Do1], we have

$$\widehat{u}_1(x, x) = \frac{\tau_0}{6} - \frac{1}{2}T_{kjk,j} - \frac{1}{4}T_{kjk}T_{ljl}, \quad (3.25)$$

$$\begin{aligned} \widehat{b}_1(T, a) = |\det B| \left\{ \widehat{C} + \text{tr}[\wedge^p \tilde{A}] \left(\frac{\tau_0}{6} - \frac{1}{2}T_{kjk,j} - \frac{1}{4}T_{kjk}T_{ljl} + \frac{1}{6}\rho_{kk} \right. \right. \\ \left. \left. + \frac{1}{3}R_{iksh}B_{ki}B_{hs} + \frac{1}{3}R_{ikth}B_{kt}B_{hi} - R_{k\alpha h\alpha}B_{ks}B_{hs} \right) \right\}. \end{aligned} \quad (3.26)$$

$$\frac{\partial \overline{W}_{l_{m_1} i_{m_1}}}{\partial y_\delta} \Big|_{y=0} = \frac{1}{2p} A_{i_{m_1} l_{m_1}} Q_{ii\delta+n}; \quad (3.27)$$

$$\begin{aligned} \frac{\partial^2 \overline{W}_{l_{m_3} i_{m_3}}}{\partial y_\delta^2} \Big|_{y=0} = A_{\varepsilon_1 l_{m_3}} A_{rl} B_{l\delta} B_{v\delta} A_{m\varepsilon_1} R_{rvmi_{m_3}} + A_{i_{m_3} l_{m_3}} \left[-\frac{1}{p} \sum_{j=n+1}^d Q_{ii\delta+n,j} B_{j\delta+n} \right. \\ \left. + \frac{1}{4p^2} Q_{ii\delta+n} Q_{kk\delta+n} + \frac{1}{2p} Q_{ii\delta+n,\delta+n} \right]. \end{aligned} \quad (3.28).$$

So

$$\begin{aligned} \widehat{C} = \sum_{\delta=1}^{d-n} \sum_{1 \leq i_1 < \dots < i_p \leq d} \sum_{l_1, \dots, l_p} \varepsilon_{i_1, \dots, i_p}^{l_1, \dots, l_p} \left[\frac{C_p^2}{2p^2} \sum_{1 \leq m_1 < m_2 \leq p} A_{l_1 i_1} \dots A_{l_p i_p} Q_{ii\delta+n} Q_{kk\delta+n} \right. \\ \left. + \sum_{1 \leq m_3 \leq p} A_{l_1 i_1} \dots \widehat{A}_{l_{m_3} i_{m_3}} \dots A_{l_p i_p} \cdot A_{\varepsilon_1 l_{m_3}} [A_{rl} B_{l\delta} B_{v\delta} A_{m\varepsilon_1} R_{rvmi_{m_3}} \right. \\ \left. + A_{l_1 i_1} \dots A_{l_p i_p} \left(- \sum_{j=n+1}^d Q_{ii\delta+n,j} B_{j\delta+n} + \frac{1}{4p} Q_{ii\delta+n} Q_{kk\delta+n} + \frac{1}{2} Q_{ii\delta+n,\delta+n} \right) \right]. \end{aligned} \quad (3.29).$$

Theorem 4 *The coefficient $\widehat{b}_k(T, a)$ is of the form $\widehat{b}_k(T, a) = |\det B| \widehat{b}'_k(T, a)$ where $\widehat{b}'_k(T, a)$ is an invariant polynomial in the components of A , B and the curvature tensor R and the torsion tensor $\overline{T}$ and its covariant derivative at a . In particular, $\widehat{b}_1(T, a)$ are determined by (3.26) and (3.29).*

4 The equivariant Gilkey-Branson-Fulling formula

Since T is a preserving orientation isometry, then T^* commutes with d , δ and Δ , so T^* preserves the Hodge decomposition. Let

$$\Delta_d^{(p)} = \Delta^{(p)}|_{\text{Im}d}; \quad \Delta_\delta^{(p)} = \Delta^{(p)}|_{\text{Im}\delta}$$

and for each $t > 0$

$$f_T(t, d^{(p)}) = \text{Tr}(T^* e^{-t\Delta_d^{(p)}}); \quad f_T(t, \delta^{(p)}) = \text{Tr}(T^* e^{-t\Delta_\delta^{(p)}}),$$

$$f_T(t, \Delta^{(p)}) = \text{Tr}(T^* e^{-t\Delta^{(p)}}); \quad \beta_{T,p} = \text{Tr}(T^*|_{\ker(\Delta^{(p)})}).$$

Then we have via the Hodge decomposition theorem:

$$f_T(t, \Delta^{(p)}) = \beta_{T,p} + f_T(t, d^{(p)}) + f_T(t, \delta^{(p)}). \quad (4.1)$$

Similar to the nonequivariant case, we have

$$f_T(t, d^{(p)}) = f_T(t, \delta^{(p-1)}). \quad (4.2)$$

By (4.1) and (4.2), we have

$$f_T(t, \delta^{(p)}) = \sum_{j \leq p} (-1)^{p-j} [f_T(t, \Delta^{(j)}) - \beta_{T,j}].$$

On the other hand,

$$f_T(t, d^{(p)}) = f_T(t, \delta^{(p-1)}) = \sum_{j \leq p-1} (-1)^{p-j} [f_T(t, \Delta^{(j)}) - \beta_{T,j}].$$

Let $D^{(p)} = a^2 d\delta + b^2 \delta d$ acting on $\wedge^p$ where $a \neq 0$, $b \neq 0$. If we make the same computations for operator $D^{(p)}$ we obtain

$$\begin{aligned} f_T(t, D^{(p)}) &= \text{Tr}(T^* e^{-tD^{(p)}}) = \beta_{T,p} + f_T(a^2 t, d^{(p)}) + f_T(b^2 t, \delta^{(p)}) \\ &= f_T(b^2 t, \Delta^{(p)}) + \sum_{j < p} (-1)^{p-j} [f_T(b^2 t, \Delta^{(j)}) - f_T(a^2 t, \Delta^{(j)})]. \end{aligned} \quad (4.3)$$

By Lemma 1.8.2 in [Gil], we have

$$\begin{aligned} f_T(t, D^{(p)}) &= \sum_{N \in \Omega} (4\pi t)^{-\frac{n_N}{2}} \sum_{k=0}^{\infty} t^k b_{N,k}^T(D^{(p)}) \\ f_T(b^2 t, \Delta^{(j)}) &= \sum_{N \in \Omega} (4\pi t)^{-\frac{n_N}{2}} b^{-n_N} \sum_{k=0}^{\infty} t^k b_{N,k}^T(\Delta^{(j)}) b^{2k} \\ f_T(a^2 t, \Delta^{(j)}) &= \sum_{N \in \Omega} (4\pi t)^{-\frac{n_N}{2}} a^{-n_N} \sum_{k=0}^{\infty} t^k b_{N,k}^T(\Delta^{(j)}) a^{2k} \end{aligned}$$

By equating coefficients of t^l in the asymptotic expansion in (4.3), we get

Theorem 5

$$\begin{aligned} \sum_{N \in \Omega} (4\pi)^{-\frac{n_N}{2}} b_{N,l+\frac{n_N}{2}}^T(D^{(p)}) &= b^{2l} \sum_{N \in \Omega} (4\pi)^{-\frac{n_N}{2}} b_{N,l+\frac{n_N}{2}}^T(\Delta^{(p)}) \\ &+ \sum_{j < p} (-1)^{p-j} (b^{2l} - a^{2l}) \sum_{N \in \Omega} (4\pi)^{-\frac{n_N}{2}} b_{N,l+\frac{n_N}{2}}^T(\Delta^{(j)}). \end{aligned} \quad (4.4)$$

Remark 1. If $n_N = \text{constant}$ and taking $l = -\frac{n_N}{2}$, then we have

$$\sum_{N \in \Omega} b_{N,0}^T(D^{(p)}) = b^{-n} \sum_{N \in \Omega} b_{N,0}^T(\Delta^{(p)}) + \sum_{j < p} (-1)^{p-j} (b^{-n} - a^{-n}) \sum_{N \in \Omega} b_{N,0}^T(\Delta^{(j)}). \quad (4.5)$$

If $n_N = \text{constant}$ and taking $l = -\frac{n_N}{2} + 1$, then we have

$$\sum_{N \in \Omega} b_{N,1}^T(D^{(p)}) = b^{-n+2} \sum_{N \in \Omega} b_{N,1}^T(\Delta^{(p)}) + \sum_{j < p} (-1)^{p-j} (b^{-n+2} - a^{-n+2}) \sum_{N \in \Omega} b_{N,1}^T(\Delta^{(j)}). \quad (4.6)$$

Remark 2. Let M be a Kähler manifold and T preserve the orientation and the canonical almost complex structure, then T^* commutes with $\partial, \bar{\partial}, \partial^*, \bar{\partial}^*$. Similar to Theorem 3, we can get the equivariant version of the expression of heat kernel coefficients of nonminimal operators on Kähler manifolds in [AV].

Acknowledgement This work was supported by NSFC No. 10801027.

References

- [AV] S. Alexandrov, D. Vassilevich, *Heat kernel for nonminimal operators on a Kahler manifold*. J. Math. Phys. 37 (1996), no. 11, 5715–5718.
- [BGV] N. Berline; E. Getzler; M. Vergne, *Heat kernels and Dirac operators*. Corrected reprint of the 1992 original. Grundlehren Text Editions. Springer-Verlag, Berlin, 2004.
- [Do1] H. Donnelly, *Heat equation asymptotics with torsion*, Indiana Univ. Math. J. 34 (1985), no. 1, 105–113.
- [Do2] H. Donnelly, *Spectrum and the fixed point sets of isometries I*. Math. Ann. 224 (1976), no. 2, 161–170.
- [DGGW] E. Dryden; C. Gordon; S. Greenwald; D. Webb, *Asymptotic expansion of the heat kernel for orbifolds*. Michigan Math. J. 56 (2008), no. 1, 205–238.
- [LYZ] J. D. Lafferty, Y. L. Yu and W. P. Zhang, *A direct geometric proof of Lefschetz fixed point formulas*, Trans. AMS. 329(1992), 571–583.
- [G] P. Gilkey, *Heat content asymptotics of nonminimal operators*. Topol. Methods Nonlinear Anal. 3 (1994), no. 1, 69–80.
- [G1] P. Gilkey, *Invariance theory, the heat equation, and the Atiyah-Singer index theorem*. Second edition. Studies in Advanced Mathematics. CRC Press, Boca Raton, FL, 1995.

- [GBF] P. Gilkey; T. Branson; S. Fulling, *Heat equation asymptotics of “nonminimal” operators on differential forms*. J. Math. Phys. 32 (1991), no. 8, 2089–2091.
- [PC] M. Puta; F. Cret, *A generalization of the Gilkey-Branson-Fulling formula*. Proceedings of the Workshop on Global Analysis, Differential Geometry, Lie Algebras (Thessaloniki, 1997), 79–82, BSG Proc., 5, Geom. Balkan Press, Bucharest, 2001.
- [Yu] Y. Yu, *Trigonometry. II*, Acta Math. Sinica (N.S.) 6 (1990), no. 1, 80–86.

School of Mathematics and Statistics , Northeast Normal University, Changchun, Jilin 130024, China ;

E-mail: *wangy581@nenu.edu.cn*